

Microdermabrasion Vancouver

Microdermabrasion Vancouver - The condition of Dermatitis Herpetiformis is a chronic skin condition characterized by blisters filled with a watery fluid. It is likewise called DH or Duhring's Disease. The name implies the meaning in that it is a skin disease that has the particular appearance like herpes, however, the condition is not related to or caused by the herpes virus. DH was initially described in 1884 by Dr. Louis Duhring, thus the name. In the year 1867, a correlation between gluten intolerance or celiac disease and DH was recognized. In nearly all of cases, the age of onset is normally 15 to 40 years, although, DH can likewise have an effect on kids as well as the elderly. Women and men are also affected. The prevalence of DH varies from 1 in 400 to 1 in 10,000.

Symptoms

Dermatitis Herpetiformis has a characteristic rash which contains chronic papulovesicular eruptions, which are extremely itchy. This particular type of rash is normally distributed symmetrically on extensor surfaces like for instance the knees, back of neck, elbows, back and the buttocks. The blisters could range from being up to 1 cm across to extremely small. Because this condition is extremely itchy, the desire to scratch can be overwhelming. This may cause the sufferer to really scratch blisters off before they are able to be checked by a medical doctor. Intense burning sensations or intense itching are occasionally felt prior to blisters appearing in a particular place.

The severity of Dermatitis Herpetiformis can vary in time if left untreated. Quite often, the initial signs appear in early adulthood, between the ages of 20 and 30 years old. The very first obvious symptoms are the small vesicles or papules that resemble bumps or blisters. The first indications are intensive burning and itching. Normally, the first blisters appear on the lower end of the spinal column, on the shoulders, along the hairline and sometimes in the mouth. Rarely, does the rash happen on several mucous membranes other than the mouth or the lips. If gluten ingestion is avoided and the correct medication is provided, the signs often disappear. In several cases, taking oral contraceptives can worsen the symptoms.

The signs of DH tend to come and go, in short periods of time, yet are always persistent. These signs can be accompanied by fatigue and abdominal pain and other indications of celiac disease.

The DH rash normally forms and disappears in 3 stages. The patient would initially see a slight discoloration of the skin where the lesions appear. The next stage would be the lesions transforming into groups of obvious vesicles and papules. The healing stage of the lesions is the final stage of development. Often this set of signs is characterized by a skin color change. This might result in parts of the skin turning darker or lighter compared to the color of the skin elsewhere on the body. Because of the intense itching, individuals normally scratch that could cause the formation of crusts.